

REPRESENTATION OF PERIOD DOUBLING BY DIGRAPHS AND CHARACTERISTIC POLYNOMIALS

RICHELL O. CELESTE
Institute of Mathematics
University of the Philippines
Diliman Quezon City
e-mail: ching@math.upd.edu.ph

YOSHIFUMI TAKENOUCHI
Institute of Mathematics
University of the Philippines
Diliman Quezon City
e-mail: jytakenovich@math.upd.edu.ph

Abstract

A general procedure which defines a partial ordering of cyclic permutations induced by continuous maps is known for constructing immediate successors to a cycle. We expound on this procedure in terms of labelled digraphs and characteristic polynomials then apply this study to period doubling, the most common route to chaos for a nonlinear dynamical system.

Key words: forcing relation; digraph; spectrum; period doubling.

1 Introduction

Li and Yorke [10] showed that the presence of a periodic orbit with least period 3 forces the presence of periodic orbit with all possible least periods. However before [10], Sharkovsky [6, 9] had already obtained the results which include the statement above as a corollary. Baldwin [3] extended this idea to get a partial order on the set of cycles in which the type of the periodic orbit is taken into account. Bernhardt [5] showed how to construct immediate successors to a given cycle.

First we reexamine [5] to obtain graph theoretical explanation of the relation between a cycle and its immediate successors. With this view point, we then obtain the representation of the period-doubling bifurcation phenomena by the digraphs

consisting of signed vertices and directed edges and by the characteristic polynomials of the adjacency matrices. Obtaining a discrete expression of the period-doubling bifurcation phenomena implies that we may define the period-doubling bifurcation phenomena on the graph itself without referring to the original one-dimensional continuous maps.

In section 2 we present basic definitions and concepts on the forcing relation on cycles in dimension one and on graph theory. In section 3 we obtain the relation between the digraph and the characteristic polynomial of any cycle and that of its immediate successor. In section 4 we describe the period-doubling phenomena of the family of the logistic curves with their digraphs and characteristic polynomials.

2 Preliminaries

Definition 2.1. Let $f : I \rightarrow I$ be a continuous map of a compact interval I to itself. We define $f^0(x) = x$ and for $n \in \mathbb{N} \setminus \{0\}$, $f^n(x) = f(f^{n-1}(x))$. If there exists $k \in \mathbb{N}$ such that $f^k(x) = x$, then we say x is **periodic** for f and x has (least) **period** s , where s is smallest element of $\mathbb{N}$ such that $f^s(x) = x$. For $x \in I$, the **orbit** of x is the set $\{f^n(x) | n \geq 0\}$. If x is periodic with least period s , then the orbit of x is the finite set $\mathcal{O}(x) = \{x, f(x), f^2(x), \dots, f^{s-1}(x)\}$.

Definition 2.2. A **cycle** (cyclic permutation) of length n is a bijection $\theta : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$ such that $\theta^k(1) \neq 1$ for $1 \leq k < n$. We denote a cycle θ by $\theta = (k_1, k_2, \dots, k_m)$, where $\theta(k_i) = k_{i+1}$ and $\theta(k_m) = k_1$. We assume, without loss of generality, that $k_1 = 1$. Write the elements of a periodic orbit $\mathcal{O}$ in increasing order: $x_1 < x_2 < \dots < x_s$. We say $\mathcal{O}$ has **orbit type** θ if θ is a cycle of length s and for each $i \in \{1, 2, \dots, s\}$, $f(x_i) = x_{\theta(i)}$.

$(S_n, \circ)$ denotes the group of permutations on n objects, C_n denotes the subset of S_n consisting of cycles (cyclic permutations) of length n . We also denote $\bigcup_{n \geq 1} C_n$ by C .

Definition 2.3. The **forcing relation on cycles** is defined as follows: $\beta \in C$ forces $\alpha \in C$ if and only if every continuous map of the interval that has a periodic orbit of type β has a periodic orbit of type α .

Baldwin [3] showed that this relation induces a partial order on the set of cycles and provided an algorithm to decide when one cycle forces another.

Definition 2.4. [5] We say $\beta \in C$ is an **immediate successor** to $\alpha \in C$, or α is an **immediate predecessor** to β , if $\beta \neq \alpha$ and the set of all cycles forced by β is same as the set of all cycles forced by α and α itself.

Definition 2.5. For $\theta \in S_n$, the **canonical θ -linear map**, or the **connect-the-dots map of θ** is defined by $L_\theta : [1, n] \rightarrow [1, n]$ where $L_\theta = \theta$ on $\{1, 2, \dots, n\}$ and L_θ is linear on $I_i := [i, i+1]$ for $i \in \{1, 2, \dots, n-1\}$.

It is known that $\beta \in C$ forces $\alpha \in C$ if and only if L_β has a periodic orbit of cycle type α [2].

Definition 2.6. For $\theta \in C_n$, $n \geq 3$ we say θ is **$+k$ -modal cycle** or **$-k$ -modal cycle** if the number of all the local extrema of L_θ is equal to k and $\theta(1) < \theta(2)$ or $\theta(1) > \theta(2)$, respectively. In particular $+1$ -modal cycle is called **unimodal cycle**.

The forcing relation defined by Baldwin induces a total order on the set of unimodal cycles [7].

Definition 2.7. From L_θ , we define the **labelled digraph (directed graph) of θ** , denoted by $G(\theta)$, as follows: $G(\theta) = (\mathcal{V}, \mathcal{E})$ consists of the set of $n-1$ signed vertices $\mathcal{V} = \{v_1^\pm, v_2^\pm, \dots, v_{n-1}^\pm\}$ where we take v_i^+ if $\theta(i) < \theta(i+1)$ or v_i^- otherwise, and the set of directed edges $\mathcal{E} = \{e_{ij}\}$ where e_{ij} a directed edge from v_i to v_j exists if $I_j \subset L_\theta(I_i)$.

Conversely, from $G(\theta)$ we can obtain L_θ . In this sense, we say they are equivalent. Indeed Baldwin's algorithm [3] which decides whether one cycle θ forces another or not, is derived from $G(\theta)$.

Definition 2.8. The **induced matrix of θ** or the **adjacency matrix of $G(\theta)$** [8], denoted by $M(\theta) = [a_{i,j}]$, is the $(n-1) \times (n-1)$ matrix with (i, j) -th entry given by

$$a_{i,j} = \begin{cases} 1, & \text{if } I_j \subset L_\theta(I_i), \\ 0, & \text{otherwise,} \end{cases}$$

where $i, j \in [1, n-1]$.

Definition 2.9. [8] A graph $H = (\mathcal{W}, \mathcal{F})$ is said to be a **subgraph** of the graph $G = (\mathcal{V}, \mathcal{E})$ if $\mathcal{W} \subset \mathcal{V}$ and $\mathcal{F} \subset \mathcal{E}$. In particular if $\mathcal{F}$ consists of all the edges from $\mathcal{E}$ which connect the vertices from $\mathcal{W}$, then H is called an **induced subgraph**.

Definition 2.10. [8] A **cycle of length n** in the digraph G is a subgraph with the vertex set $\{v_1, \dots, v_n\}$ having directed edges from v_i to v_{i+1} where $i = 1, \dots, n-1$ and a directed edge from v_n to v_1 . In particular if $v_1 < v_2 < \dots < v_n$ then this cycle induces the cyclic permutation $\theta = (12 \dots n)$.

3 Theorem

Definition 3.1. [5] Let $\theta \in C_n$, then $\theta^* \in S_{2n}$ is defined by

$$\theta^*(2k) := 2\theta(k), \quad \theta^*(2k-1) := 2\theta(k) - 1,$$

where $k \in \{1, 2, \dots, n\}$.

Definition 3.2. [5] Let ρ_s denote the transposition $(2s-1, 2s)$, where $s \in \mathbb{N}$. Then for $\theta \in C_n$ define η as follows:

$$\eta := \theta^* \circ \underbrace{\rho_{i_1} \circ \rho_{i_2} \circ \dots \circ \rho_{i_m}}_m,$$

where $1 \leq i_1 < i_2 < \dots < i_m \leq n$.

If m is odd, then $\eta \in C_{2n}$ and it is an immediate successor to θ . If m is even, then $\eta \in S_{2n} \setminus C_{2n}$. There are 2^{n-1} immediate successors $\eta \in C_{2n}$ to θ , and 2^{n-1} non-cyclic permutations $\eta \in S_{2n}$. (See [4] and [5])

For θ and η above, denote their labelled digraphs by $G(\theta)$ and $G(\eta)$, respectively. Then $G(\theta) = (\mathcal{V}, \mathcal{E})$ consists of the set of $n-1$ signed vertices $\mathcal{V} = \{v_1^\pm, v_2^\pm, \dots, v_{n-1}^\pm\}$, where we take v_i^+ if $\theta(i) < \theta(i+1)$ or v_i^- otherwise, and the set of directed edges $\mathcal{E} = \{e_{ij}\}$ where e_{ij} a directed edge from v_i to v_j exists if $I_j \subset L_\theta(I_i)$. $G(\eta) = (\mathcal{W}, \mathcal{F})$ consists of the set of $2n-1$ signed vertices $\mathcal{W} = \{w_1^\pm, w_2^\pm, \dots, w_{2n-1}^\pm\}$, where we take w_i^+ if $\eta(i) < \eta(i+1)$ or w_i^- otherwise, and the set of directed edges $\mathcal{F} = \{f_{ij}\}$ where f_{ij} a directed edge from w_i to w_j exists if $I_j \subset L_\eta(I_i)$. Let us divide $\mathcal{W}$ into two subsets $\mathcal{W}_1 = \{w_2^\pm, \dots, w_{2n-2}^\pm\}$ and $\mathcal{W}_2 = \{w_1^\pm, w_3^\pm, \dots, w_{2n-1}^\pm\}$.

Then we have the following observations:


(i) $G(\eta)|_{\mathcal{W}_1}$, the induced subgraph of $G(\eta) = (\mathcal{W}, \mathcal{F})$ determined on $\mathcal{W}_1$ is isomorphic to the digraph $G(\theta) = (\mathcal{V}, \mathcal{E})$. Indeed, the isomorphism $\varphi : \mathcal{V} \rightarrow \mathcal{W}_1$ is given by

$$\varphi(v_i^\pm) = w_{2i}^\pm$$

where $i \in \{1, 2, \dots, n-1\}$. Observe that the sign of vertex v_i is always consistent with that of vertex w_{2i} .

(ii) In $G(\eta) = (\mathcal{W}, \mathcal{F})$ there does not exist any directed edge from $\mathcal{W}_2$ to $\mathcal{W}_1$.

(iii) $G(\eta)|_{\mathcal{W}_2}$, the induced subgraph of $G(\eta) = (\mathcal{W}, \mathcal{F})$ determined on $\mathcal{W}_2$, consists of an n -cycle induced by θ . Indeed, this n -cycle is given by



and the type of this n -cycle is also θ .

Together, (i), (ii), and (iii) provide a graph theoretical explanation of the reason why η is an immediate successor to θ (see LEMMA 2.4 in [5]). As a consequence, we have the following theorem.

Theorem 3.1. *Denote the characteristic polynomials of the adjacency matrices of $G(\theta)$, $G(\eta)$, $G(\eta)|_{\mathcal{W}_1}$ and $G(\eta)|_{\mathcal{W}_2}$ by $P_\theta(\lambda)$, $P_\eta(\lambda)$, $P_{\mathcal{W}_1}(\lambda)$ and $P_{\mathcal{W}_2}(\lambda)$, respectively. Then $P_\eta(\lambda)$ is given by the product of $P_{\mathcal{W}_1}(\lambda)$ and $P_{\mathcal{W}_2}(\lambda)$, namely*

$$P_\eta(\lambda) = P_\theta(\lambda) \cdot (\lambda^n - 1).$$

As a corollary of this theorem, we have the following.

Corollary 3.1. *The spectrum of $G(\eta)$ consists of the spectrum of $G(\theta)$ and the n distinct roots of unity.*

Assume that the adjacency matrix of $G(\theta)$ is given by

$$M(\theta) = \begin{bmatrix} a_{i,j} \end{bmatrix}_{(n-1) \times (n-1)},$$

where $1 \leq i, j \leq n-1$, and the adjacency matrix of $G(\eta)$ is given by

$$M(\eta) = \begin{bmatrix} b_{i,j} \end{bmatrix}_{(2n-1) \times (2n-1)},$$

where $1 \leq i, j \leq 2n-1$.

Then we have the following Lemma 3.1, and 3.2.

Lemma 3.1. *$G(\eta)|_{\mathcal{W}_1}$ is isomorphic to $G(\theta)$ by φ , namely*

$$b_{2i,2j} = a_{i,j},$$

where $1 \leq i, j \leq n-1$.

Proof. For any $\theta \in C_n$, $\theta^* \in S_{2n}$ is defined by

$$\theta^* := \begin{pmatrix} \cdots & 2i-1 & 2i & 2i+1 & 2i+2 & \cdots \\ \cdots & 2\theta(i)-1 & 2\theta(i) & 2\theta(i+1)-1 & 2\theta(i+1) & \cdots \end{pmatrix}.$$

If $\eta := \theta^* \circ \cdots \circ \rho_i \circ \cdots$, then

$$\eta = \begin{pmatrix} \cdots & 2i-1 & 2i & 2i+1 & 2i+2 & \cdots \\ \cdots & 2\theta(i) & 2\theta(i)-1 & 2\theta(i+1)-1 & 2\theta(i+1) & \cdots \end{pmatrix},$$

and

$$L_\eta([2i, 2i+1]) = \begin{cases} [2\theta(i)-1, 2\theta(i+1)-1] & \text{if } \theta(i) < \theta(i+1) \\ [2\theta(i+1)-1, 2\theta(i)-1] & \text{if } \theta(i) > \theta(i+1) \end{cases}.$$

If $\eta := \theta^* \circ \cdots \circ \rho_{i+1} \circ \cdots$, then

$$\eta = \begin{pmatrix} \cdots & 2i-1 & 2i & 2i+1 & 2i+2 & \cdots \\ \cdots & 2\theta(i)-1 & 2\theta(i) & 2\theta(i+1) & 2\theta(i+1)-1 & \cdots \end{pmatrix},$$

and

$$L_\eta([2i, 2i+1]) = \begin{cases} [2\theta(i), 2\theta(i+1)] & \text{if } \theta(i) < \theta(i+1) \\ [2\theta(i+1), 2\theta(i)] & \text{if } \theta(i) > \theta(i+1) \end{cases}.$$

If $\eta := \theta^* \circ \cdots \circ \rho_i \circ \rho_{i+1} \circ \cdots$, then

$$\eta = \begin{pmatrix} \cdots & 2i-1 & 2i & 2i+1 & 2i+2 & \cdots \\ \cdots & 2\theta(i) & 2\theta(i)-1 & 2\theta(i+1) & 2\theta(i+1)-1 & \cdots \end{pmatrix},$$

and

$$L_\eta([2i, 2i+1]) = \begin{cases} [2\theta(i)-1, 2\theta(i+1)] & \text{if } \theta(i) < \theta(i+1) \\ [2\theta(i+1), 2\theta(i)-1] & \text{if } \theta(i) > \theta(i+1) \end{cases}.$$

If $\eta := \theta^* \circ \rho_{i_1} \circ \rho_{i_2} \circ \cdots \circ \rho_{i_m}$ does not include ρ_i nor ρ_{i+1} , then

$$\eta = \begin{pmatrix} \cdots & 2i-1 & 2i & 2i+1 & 2i+2 & \cdots \\ \cdots & 2\theta(i)-1 & 2\theta(i) & 2\theta(i+1)-1 & 2\theta(i+1) & \cdots \end{pmatrix},$$

and

$$L_\eta([2i, 2i+1]) = \begin{cases} [2\theta(i), 2\theta(i+1)-1] & \text{if } \theta(i) < \theta(i+1) \\ [2\theta(i+1)-1, 2\theta(i)] & \text{if } \theta(i) > \theta(i+1) \end{cases}.$$

Under the assumption $\theta(i) < \theta(i+1)$, we have

$$\begin{aligned} & a_{i,j} = 1 \\ \Leftrightarrow & I_j \subset L_\theta(I_i) \\ \Leftrightarrow & [j, j+1] \subset [\theta(i), \theta(i+1)] \\ \Leftrightarrow & \theta(i) \leq j \text{ and } j+1 \leq \theta(i+1) \\ \Leftrightarrow & 2\theta(i) \leq 2j \text{ and } 2j+1 \leq 2\theta(i+1)-1 \\ \Leftrightarrow & [2j, 2j+1] \subset [2\theta(i), 2\theta(i+1)-1]. \end{aligned}$$

Here we note that

$$[2\theta(i), 2\theta(i+1) - 1] \subset [2\theta(i) - 1, 2\theta(i+1) - 1], [2\theta(i), 2\theta(i+1)] \subset [2\theta(i) - 1, 2\theta(i+1)],$$

and

$$\underset{\text{even}}{[2j, 2j+1]} \neq \underset{\text{odd}}{[2\theta(i) - 1, 2\theta(i)]}, \underset{\text{odd}}{[2\theta(i+1) - 1, 2\theta(i+1)]}, \underset{\text{even}}{[2\theta(i), 2\theta(i+1)]},$$

thus,

$$\begin{aligned} & [2j, 2j+1] \subset [2\theta(i), 2\theta(i+1) - 1] \\ \Leftrightarrow & [2j, 2j+1] \subset [\eta(2i), \eta(2i+1)] \\ \Leftrightarrow & I_{2j} \subset L_\eta(I_{2i}) \\ \Leftrightarrow & b_{2i, 2j} = 1. \end{aligned}$$

Similarly, under the assumption $\theta(i) > \theta(i+1)$, we can also show that

$$a_{i,j} = 1 \Leftrightarrow b_{2i, 2j} = 1.$$

This concludes the proof of the lemma. $\square$

Lemma 3.2. *In $G(\eta) = (\mathcal{W}, \mathcal{F})$, there does not exist any directed edge from $\mathcal{W}_2$ to $\mathcal{W}_1$, that is*

$$b_{2i-1, 2j} = 0,$$

where $1 \leq i \leq n$, $1 \leq j \leq n-1$. Moreover, $G(\eta)|_{\mathcal{W}_2}$ consists of an n -cycle induced by θ , namely

$$\begin{cases} b_{2i-1, 2j-1} = 1 & \text{if } j = \theta(i), \\ b_{2i-1, 2j-1} = 0 & \text{otherwise,} \end{cases}$$

where $1 \leq i, j \leq n$.

In other words

$$\begin{bmatrix} b_{12} & b_{14} & \cdots & b_{1(2n-2)} \\ b_{32} & b_{34} & \cdots & b_{3(2n-2)} \\ \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ b_{(2n-1)2} & b_{(2n-1)4} & \cdots & b_{(2n-1)(2n-2)} \end{bmatrix} = O_{n \times (n-1)},$$

and

$$\begin{bmatrix} b_{11} & b_{13} & \cdots & b_{1(2n-1)} \\ b_{31} & b_{33} & \cdots & b_{3(2n-1)} \\ \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ b_{(2n-1)1} & b_{(2n-1)3} & \cdots & b_{(2n-1)(2n-1)} \end{bmatrix} = \begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1n} \\ c_{21} & c_{22} & \cdots & c_{2n} \\ \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ c_{n1} & c_{n2} & \cdots & c_{nn} \end{bmatrix}$$

$$\Longleftrightarrow b_{2i-1,2j-1} = c_{ij},$$

where

$$\begin{cases} c_{ij} = 1 & \text{if } j = \theta(i), \\ c_{ij} = 0 & \text{otherwise.} \end{cases}$$

Proof.

$$\begin{aligned} & j = \theta(i) \\ \Leftrightarrow & \theta^*(2i-1) = 2j-1 \text{ and } \theta^*(2i) = 2j \\ \Leftrightarrow & [2j-1, 2j] = [\theta^*(2i-1), \theta^*(2i)]. \end{aligned}$$

For any $\theta \in C_n$, $\theta^* \in S_{2n}$ is defined by

$$\theta^* := \begin{pmatrix} \cdots & 2i-1 & 2i & \cdots \\ \cdots & \theta^*(2i-1) & \theta^*(2i) & \cdots \end{pmatrix}.$$

If $\eta := \theta^* \circ \cdots \circ \rho_i \circ \cdots$, then

$$\eta = \begin{pmatrix} \cdots & 2i-1 & 2i & \cdots \\ \cdots & \theta^*(2i) & \theta^*(2i-1) & \cdots \end{pmatrix},$$

and thus $[2j-1, 2j] = [\theta^*(2i-1), \theta^*(2i)] = [\eta(2i), \eta(2i-1)]$.

If $\eta := \theta^* \circ \rho_{i_1} \circ \rho_{i_2} \circ \cdots \circ \rho_{i_m}$ does not include ρ_i , then

$$\eta = \begin{pmatrix} \cdots & 2i-1 & 2i & \cdots \\ \cdots & \theta^*(2i-1) & \theta^*(2i) & \cdots \end{pmatrix},$$

and thus $[2j-1, 2j] = [\theta^*(2i-1), \theta^*(2i)] = [\eta(2i-1), \eta(2i)]$.

In either event, we have

$$\begin{aligned} & [2j-1, 2j] = [\theta^*(2i-1), \theta^*(2i)] \\ \Leftrightarrow & I_{2j-1} = L_\eta(I_{2i-1}) \\ \Leftrightarrow & \begin{cases} b_{2i-1,2j-1} = 1 \\ b_{2i-1,k} = 0 \end{cases} \quad \text{if } k \neq 2j-1, \end{aligned}$$

ending the proof of the lemma. □

The following statement is obvious:

Lemma 3.3.

$$\begin{vmatrix} A_{l \times l} & O_{l \times m} \\ C_{m \times l} & B_{m \times m} \end{vmatrix} = |A_{l \times l}| |B_{m \times m}|.$$

We are now ready to prove the theorem.

Proof of Theorem 3.1. The characteristic polynomial of the adjacency matrix of $G(\theta)$ is given by

$$P_\theta(\lambda) = |\lambda I - M(\theta)| = \begin{vmatrix} \lambda - a_{11} & -a_{12} & \cdots & -a_{1(n-1)} \\ -a_{21} & \lambda - a_{22} & \cdots & -a_{2(n-1)} \\ \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots \\ -a_{(n-1)1} & -a_{(n-1)2} & \cdots & \lambda - a_{(n-1)(n-1)} \end{vmatrix}.$$

Let

$$\sigma_{2n-1} := \begin{pmatrix} 1 & 2 & 3 & \cdots & n & n+1 & n+2 & n+3 & \cdots & 2n-1 \\ 1 & 3 & 5 & \cdots & 2n-1 & 2 & 4 & 6 & \cdots & 2n-2 \end{pmatrix}.$$

Using Lemmas 3.1, 3.2, and 3.3, the characteristic polynomial $P_\eta(\lambda)$ of η is thus

$$\begin{aligned} &= \begin{vmatrix} \lambda - b_{11} & -b_{12} & \cdots & \cdots & -b_{1(2n-2)} & -b_{1(2n-1)} \\ -b_{21} & \lambda - b_{22} & \cdots & \cdots & -b_{2(2n-2)} & -b_{2(2n-1)} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -b_{(2n-2)1} & -b_{(2n-2)2} & \cdots & \cdots & \lambda - b_{(2n-2)(2n-2)} & -b_{(2n-2)(2n-1)} \\ -b_{(2n-1)1} & -b_{(2n-1)2} & \cdots & \cdots & -b_{(2n-1)(2n-2)} & \lambda - b_{(2n-1)(2n-1)} \end{vmatrix} \\ &= \text{sgn}(\sigma_{2n-1}) \begin{vmatrix} \lambda - b_{11} & -b_{12} & \cdots & \cdots & -b_{1(2n-2)} & -b_{1(2n-1)} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -b_{(2n-2)1} & -b_{(2n-2)2} & \cdots & \cdots & \lambda - b_{(2n-2)(2n-2)} & -b_{(2n-2)(2n-1)} \\ -b_{21} & \lambda - b_{22} & \cdots & \cdots & -b_{2(2n-2)} & -b_{2(2n-1)} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -b_{(2n-1)1} & -b_{(2n-1)2} & \cdots & \cdots & -b_{(2n-1)(2n-2)} & \lambda - b_{(2n-1)(2n-1)} \end{vmatrix} \\ &= (\text{sgn}(\sigma_{2n-1}))^2 \begin{vmatrix} \lambda - b_{11} & \cdots & -b_{1(2n-1)} & -b_{12} & \cdots & -b_{1(2n-2)} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -b_{(2n-1)1} & \cdots & \lambda - b_{(2n-1)(2n-1)} & -b_{(2n-1)2} & \cdots & -b_{(2n-1)(2n-2)} \\ -b_{21} & \cdots & -b_{2(2n-1)} & \lambda - b_{22} & \cdots & -b_{2(2n-2)} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -b_{(2n-2)1} & \cdots & -b_{(2n-2)(2n-1)} & -b_{(2n-2)2} & \cdots & \lambda - b_{(2n-2)(2n-2)} \end{vmatrix} \\ &= \begin{vmatrix} \lambda - b_{11} & \cdots & -b_{1(2n-1)} \\ \cdots & \cdots & \cdots \\ -b_{(2n-1)1} & \cdots & \lambda - b_{(2n-1)(2n-1)} \end{vmatrix} \begin{vmatrix} \lambda - b_{22} & \cdots & -b_{2(2n-2)} \\ \cdots & \cdots & \cdots \\ -b_{(2n-2)2} & \cdots & \lambda - b_{(2n-2)(2n-2)} \end{vmatrix} \\ &= (\lambda^n + \text{sgn}(\theta)(-1)^n)P_\theta(\lambda) \end{aligned}$$

since $(\text{sgn}(\sigma_{2n-1}))^2 = 1$,

$$\begin{bmatrix} -b_{12} & \cdots & -b_{1(2n-2)} \\ \cdots & \cdots & \cdots \\ -b_{(2n-1)2} & \cdots & -b_{(2n-1)(2n-2)} \end{bmatrix} = O_{n \times (n-1)},$$

$$\begin{vmatrix} \lambda - b_{22} & \cdots & -b_{2(2n-2)} \\ \cdots & \cdots & \cdots \\ -b_{(2n-2)2} & \cdots & \lambda - b_{(2n-2)(2n-2)} \end{vmatrix} = \begin{vmatrix} \lambda - a_{11} & \cdots & -a_{1(n-1)} \\ \cdots & \cdots & \cdots \\ -a_{(n-1)1} & \cdots & \lambda - a_{(n-1)(n-1)} \end{vmatrix} = P_\theta(\lambda),$$

and

$$\begin{aligned} & \begin{vmatrix} \lambda - b_{11} & \cdots & -b_{1(2n-1)} \\ \cdots & \cdots & \cdots \\ -b_{(2n-1)1} & \cdots & \lambda - b_{(2n-1)(2n-1)} \end{vmatrix} \\ &= \begin{vmatrix} \lambda - c_{11} & \cdots & -c_{1n} \\ \cdots & \cdots & \cdots \\ -c_{n1} & \cdots & \lambda - c_{nn} \end{vmatrix} \text{ where } \begin{cases} c_{ij} = 1 & \text{if } j = \theta(i), \\ c_{ij} = 0 & \text{otherwise.} \end{cases} \\ &= \text{sgn}(\iota) \cdot (\lambda - 0)^n + \text{sgn}(\theta) \cdot (-1)^n \\ &= \lambda^n + \text{sgn}(\theta)(-1)^n. \end{aligned}$$

Here we note that $\text{sgn}(\theta)(-1)^n$ is always equal to -1 since

$$\text{sgn}(\theta)(-1)^n = \begin{cases} (+1) \cdot (-1) & \text{if } n \text{ is odd,} \\ (-1) \cdot (+1) & \text{if } n \text{ is even.} \end{cases}$$

This completes the proof of the theorem. $\square$

4 Immediate Successors and Period Doubling

4.1 The logistic map

The logistic map $[1, 9]$ is written as

$$x_{n+1} = ax_n(1 - x_n) \tag{1}$$

where the current value x_n , is mapped onto the next value x_{n+1} . Here, we restrict our considerations to $0 \leq x_n \leq 1$ and $0 \leq a \leq 4$. The corresponding function to the logistic map, called the logistic function or the logistic curve, is given by

$$f_a(x) = ax(1 - x) \tag{2}$$

By repeatedly iterating the logistic map forward through time, we may observe different behaviors of the iterated solutions. The sequence of iterated solutions to the map is called an orbit. The behavior of orbits, which originate from typically chosen initial condition used for the iterations, depends on the control parameter a . The logistic curve is parabolic like the quadratic function with $f_a(0) = f_a(1) = 0$ and a maximum at $x = 1/2$. Being a smooth curve with only one critical point, that is, a local maximum, it is classified as a unimodal (single humped) map.




Figure 1. Bifurcation diagram of the logistic map

4.2 Period-doubling bifurcations

The logistic map has two fixed points: 0 and $(a - 1)/a$. If $a < 1$, then $|f'_a(0)| < 1$ and we say that the fixed point $x = 0$ is stable for $a < 1$. If $a > 1$, then $|f'_a(0)| > 1$ and so the fixed point $x = 0$ is now unstable and a new stable fixed point $x = (a - 1)/a$ emerges. The fixed point $x = (a - 1)/a$ is stable for $1 < a < 3$. At $a = 3$, the fixed point $x = (a - 1)/a$ becomes unstable and a stable 2-cycle (period 2 attractor) $\{x, y\} = \{(a + 1 - \sqrt{a^2 - 2a - 3})/2a, (a + 1 + \sqrt{a^2 - 2a - 3})/2a\}$ is said to bifurcate from this fixed point. The above-mentioned 2-cycle is the solution of the system of equations given by $y = ax(1 - x)$ and $x = ay(1 - y)$. When $a = 1 + \sqrt{6} \approx 3.44949$, this 2-cycle loses its stability and a stable 4-cycle begins followed by a stable 8-cycle (between $a \approx 3.54409$ and $a \approx 3.56440$). The period continues doubling over shorter intervals until $a \approx 3.56995$ where the chaotic regime takes over. A period 3 attractor is born at $a = 1 + \sqrt{8} \approx 3.82843$ [11], causing a notably large window in the bifurcation diagram (see Figure 1). This 3-cycle also undergoes a period doubling cascade in which period $3 \cdot 2^\ell$ attractors are successively produced until chaos. This mechanism by which each $(k2^\ell)$ -cycle loses its stability and is replaced by a stable $(k2^{\ell+1})$ -cycle is called **period-doubling bifurcation** (see Figure 2).




Figure 2. Magnification of the first few period doubling


To determine period 4, period 8, and in general, period n solutions of the logistic map, we solve the system of difference equations, $x_1 = x_2, x_2 = x_3, \dots$, and $x_n = x_1$. We note that these computations become very complicated even for relatively small values of n . Finding the location of these bifurcation points on the bifurcation diagram also becomes very difficult. Even so, examining the types of orbits and their characteristic polynomials turns out to be less tortuous. This is illustrated in the following sections.

4.3 Period-doubling bifurcation beginning with nontrivial fixed point

The attracting period 2 orbit of type $\theta_1 = (12)$ bifurcates from the fixed point of type $\theta_0 = (1)$ at $a = 3$. θ_1 is the only one immediate successor to θ_0 , and it follows that

$$\begin{aligned} P_{\theta_1}(\lambda) &= \lambda - 1 \\ &= (\lambda - 1) \cdot P_{\theta_0}(\lambda), \end{aligned}$$


if we define $P_{\theta_0}(\lambda) := 1$ for $\theta_0 = (1)$. Indeed the graph $G(\theta_1) = (\mathcal{V}_1, \mathcal{E}_1)$ consists of a vertex v_{11} with the minus sign and a directed edge from v_{11} to itself (a loop associated with v_{11}):



The attracting period 4 orbit of type $\theta_2 = (1324)$ bifurcates from the period 2 orbit of type θ_1 at $a = 1 + \sqrt{6} \approx 3.44949$. θ_2 with modality $+1$ is one of $2^{2-1} = 2$ immediate successors to θ_1 . Here we note that (1423) with modality -1 is another immediate successor to θ_1 , however, the logistic map can realize only $+1$ -modal (unimodal) cycles. Now it follows that

$$\begin{aligned} P_{\theta_2}(\lambda) &= \lambda^3 - \lambda^2 - \lambda + 1 \\ &= (\lambda^2 - 1)(\lambda - 1) \\ &= (\lambda^2 - 1) \cdot P_{\theta_1}(\lambda). \end{aligned}$$

Indeed the graph $G(\theta_2) = (\mathcal{V}_2, \mathcal{E}_2)$ is given as follows:




where $\mathcal{V}_2 = \{v_{21}^+, v_{22}^-, v_{23}^-\}$, $\mathcal{V}_{21} = \{v_{22}^-\}$, and $\mathcal{V}_{22} = \{v_{21}^+, v_{23}^-\}$. Here we note that (i) $G(\theta_2)|_{\mathcal{V}_{21}}$ is isomorphic to $G(\theta_1)$, the sign of vertex v_{11} is consistent with that of vertex v_{22} , (ii) there are no directed edges from $\mathcal{V}_{22}$ to $\mathcal{V}_{21}$, (iii) $G(\theta_2)|_{\mathcal{V}_{22}}$ consists of a 2-cycle induced by θ_1 .

The attracting period 8 orbit of type $\theta_3 = (15472638)$ bifurcates from the period 4 orbit of type θ_2 at $a \approx 3.54409$. θ_3 with modality +1 is one of $2^{4-1} = 8$ immediate successors to θ_2 . There are other 7 immediate successors to θ_2 , however, their modalities are not equal to +1. Now it follows that

$$\begin{aligned}
P_{\theta_3}(\lambda) &= \lambda^7 - \lambda^6 - \lambda^5 + \lambda^4 - \lambda^3 + \lambda + \lambda - 1 \\
&= (\lambda^4 - 1)(\lambda^2 - 1)(\lambda - 1) \\
&= (\lambda^4 - 1) \cdot P_{\theta_2}(\lambda).
\end{aligned}$$

Indeed the graph $G(\theta_3) = (\mathcal{V}_3, \mathcal{E}_3)$ is given as follows:




where $\mathcal{V}_3 = \{v_{31}^+, v_{32}^+, v_{33}^-, v_{34}^-, v_{35}^-, v_{36}^-, v_{37}^-\}$, $\mathcal{V}_{31} = \{v_{32}^+, v_{34}^-, v_{36}^-\}$, and $\mathcal{V}_{32} = \{v_{31}^+, v_{33}^-, v_{35}^-, v_{37}^-\}$. Here we note that (i) $G(\theta_3)|_{\mathcal{V}_{31}}$ is isomorphic to $G(\theta_2)$, the sign of vertex $v_{3(2j)}$ is consistent with that of vertex v_{2j} ($j = 1, 2, 3$), (ii) there are no directed edges from $\mathcal{V}_{32}$ to $\mathcal{V}_{31}$, (iii) $G(\theta_3)|_{\mathcal{V}_{32}}$ consists of a 4-cycle induced by θ_2 .

Inductively, for the attracting period 2^ℓ orbit of type θ_ℓ with modality $+1$, which is one of $2^{(2^{\ell-1}-1)}$ immediate successors to $\theta_{\ell-1}$, and bifurcated from the period $2^{\ell-1}$ orbit of type $\theta_{\ell-1}$, we obtain $G(\theta_\ell)$ and $P_{\theta_\ell}(\lambda)$.

4.4 Period-doubling bifurcation beginning with period 3

Next consider the period-doubling bifurcation beginning with period 3 orbit of type $\theta_0 = (123)$. The graph $G(\theta_0) = (\mathcal{V}_0, \mathcal{E}_0)$ consists of two vertices with a 2-cycle and a loop:




and thus $P_{\theta_0}(\lambda) = \lambda^2 - \lambda - 1$.

The attracting period 6 orbit of type $\theta_1 = (135246)$ bifurcates from the period 3 orbit of type $\theta_0 = (123)$ at $a \approx 3.8415$. θ_1 with modality $+1$ is one of $2^{3-1} = 4$ immediate successors to θ_0 . (146235) , (136245) , (145236) are also immediate successors to θ_0 , however, their modalities are not equal to $+1$. Now it follows that

$$\begin{aligned} P_{\theta_1}(\lambda) &= \lambda^5 - \lambda^4 - \lambda^3 - \lambda^2 + \lambda + 1 \\ &= (\lambda^3 - 1)(\lambda^2 - \lambda - 1) \\ &= (\lambda^3 - 1) \cdot P_{\theta_0}(\lambda) \end{aligned}$$

Indeed the graph $G(\theta_1) = (\mathcal{V}_1, \mathcal{E}_1)$ is given as follows:



where $\mathcal{V}_1 = \{v_{11}^+, v_{12}^+, v_{13}^+, v_{14}^-, v_{15}^-\}$, $\mathcal{V}_{11} = \{v_{12}^+, v_{14}^-\}$, and $\mathcal{V}_{12} = \{v_{11}^+, v_{13}^+, v_{15}^-\}$. Here we note that (i) $G(\theta_1)|_{\mathcal{V}_{11}}$ is isomorphic to $G(\theta_0)$, the sign of vertex $v_{1(2j)}$ is consistent with that of vertex v_{0j} ($j = 1, 2$), (ii) there are no directed edges from $\mathcal{V}_{12}$ to $\mathcal{V}_{11}$, (iii) $G(\theta_1)|_{\mathcal{V}_{12}}$ consists of a 3-cycle induced by θ_0 .

Inductively, for the attracting period $(3 \cdot 2^\ell)$ orbit of type θ_ℓ with modality $+1$, which is one of $2^{(3 \cdot 2^{\ell-1}-1)}$ immediate successors to $\theta_{\ell-1}$, and bifurcated from the period $2^{\ell-1}$ orbit of type $\theta_{\ell-1}$, we obtain $G(\theta_\ell)$ and $P_{\theta_\ell}(\lambda)$.

4.5 Period-doubling bifurcation beginning with period k

It is known that periodic attractors exist for all odd integers k . Therefore, in general, for the attracting period $(k \cdot 2^\ell)$ orbit of type θ_ℓ with modality $+1$, which is one of $2^{(k \cdot 2^{\ell-1}-1)}$ immediate successors to $\theta_{\ell-1}$, and bifurcated from the period $2^{\ell-1}$ orbit of type $\theta_{\ell-1}$, it follows that

$$P_{\theta_\ell}(\lambda) = P_{\theta_0}(\lambda) \prod_{i=1}^{\ell} (\lambda^{(k \cdot 2^{i-1})} - 1),$$

where $P_{\theta_0}(\lambda)$ is the characteristic polynomial of the induced matrix of θ_0 , and θ_0 is the type of the original period k orbit.

Indeed the graph $G(\theta_\ell)$ consists of the set of vertices $\mathcal{V}_\ell$ and the set of directed edges $\mathcal{E}_\ell$ where $\mathcal{V}_\ell = \{v_{\ell 1}, v_{\ell 2}, v_{\ell 3}, \dots, v_{\ell(k \cdot 2^\ell - 2)}, v_{\ell(k \cdot 2^\ell - 1)}\}$, $\mathcal{V}_{\ell 1} = \{v_{\ell 2}, \dots, v_{\ell(k \cdot 2^\ell - 2)}\}$, and $\mathcal{V}_{\ell 2} = \{v_{\ell 1}, v_{\ell 3}, \dots, v_{\ell(k \cdot 2^\ell - 1)}\}$. Here we note that (i) $G(\theta_i)|_{\mathcal{V}_{i1}} \cong G(\theta_{i-1})$ for $i = 1, 2, \dots, \ell$, and the signs of the vertices are preserved by each of the above isomorphisms, (ii) there are no directed edges from $\mathcal{V}_{i2}$ to $\mathcal{V}_{i1}$ for $i = 1, 2, \dots, \ell$, (iii) $G(\theta_i)|_{\mathcal{V}_{i2}}$ consists of a $(k \cdot 2^{i-1})$ -cycle induced by θ_{i-1} for $i = 1, 2, \dots, \ell$, and $G(\theta_0)$ consists of $k - 1$ vertices and some directed edges.

In other words, $G(\theta_\ell)$ essentially consists of $k - 1$ vertices with some directed edges, a k -cycle of type θ_0 , a $2k$ -cycle of type θ_1 , $\dots$, a $(k \cdot 2^{\ell-1})$ -cycle of type $\theta_{\ell-1}$. That is the reason why $P_{\theta_\ell}(x)$ can be expressed as a product of $P_{\theta_0}(\lambda)$, $\lambda^k - 1$, $\lambda^{2k} - 1$, $\dots$, $\lambda^{(k \cdot 2^{\ell-1})} - 1$. Thus the spectrum of $G(\theta_\ell)$ consists of the spectrum of $G(\theta_0)$ and $(k \cdot 2^{i-1})$ distinct roots of $\lambda^{(k \cdot 2^{i-1})} - 1 = 0$ where $i = 1, 2, \dots, \ell$.

Remark 4.1. Conversely, we may define $G(\theta_\ell)$, the doubling of the graph $G(\theta_{i-1})$ by properties (i), (ii), and (iii).

5 Acknowledgement

The authors wish to thank Professor Evelyn L. Tan for referring to us [8].

References

- [1] Paul S. Addison, *Fractals and Chaos An Illustrated Course*, IOP Publishing Ltd (1997).

- [2] L. Alseda, J. Llibre, M. Misiurewicz, *Combinatorial Dynamics and Entropy in Dimension One*, 2nd ed., World Sci., (2000).
- [3] S. Baldwin, *Generalizations of a theorem of Sarkovskii on orbits of continuous real-valued functions*, Discrete Mathematics, Vol 67 Issue 2 (1987), 111-127.
- [4] C. Bernhardt, *Simple permutations with order a power of two*, Ergod Th. & Dynam. Sys., 4, 179 - 186 (1984).
- [5] C. Bernhardt, *The ordering on permutations induced by continuous maps of the real line*, Ergod Th. & Dynam. Sys., 7, 155 - 160 (1987).
- [6] K. Burns, B. Hasselblatt, *Sharkovsky's Theorem*, (preprint), (2007).
- [7] P. Collet and J.-P. Eckmann, *Iterated Maps on the Interval as Dynamical Systems*, Progress in Physics, vol. 1, Birkhauser, Boston, Mass., (1980)
- [8] D. M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs: Theory and Applications*, (3rd rev. enl. ed.), New York, Wiley, (1998).
- [9] P. G. Drazin, *Nonlinear systems (Cambridge Texts in Applied Mathematics)*, Cambridge University Press (1997).
- [10] T.-Y. Li, J.A. Yorke, *Period three implies Chaos*, Amer. Math. Monthly, 82 no.10 (1975), 985-992.
- [11] P. Saha, S.H. Strogaz, *The Birth of Period Three*, Mathematics Magazine, Vol. 68 No 1 Feb. (1995), 42-47.